



Please amend claim 1 as follows:

1. (Twice amended.) A pyrrolobenzodiazepine compound of the formula **Ia** or **Ib**:



(Ia)



(Ib)

wherein:

A is CH₂, or a single bond;

R₂ is selected from: R, OH, OR, CO₂H, CO₂R, COH, COR, SO₂R, CN, CH₂OR or CH=CR^AR^B, where R^A and R^B are independently selected from H, R^C, COR^C, CONH₂, CONHR^C, CONR₂^C, cyano or phosphonate, where R^C is an unsubstituted alkyl group having 1 to 4 carbon atoms;

R₆, R₇ and R₉ are independently selected from H, R, OH, OR, halo, amino, NHR, nitro, Me₃Sn;

where R is a lower alkyl group having 1 to 10 carbon atoms, or an aralkyl group of up to 12 carbon atoms, whereof the alkyl group optionally contains one or more carbon-carbon double or triple bonds, which may form part of a conjugated system, or one or more carbonyl groups or one or more ether or thioether groups, or an aryl group of up to 12 carbon atoms; and is optionally substituted by one or more halo, hydroxy, amino, or nitro groups;

and R₈ is selected from H, R, OH, OR, halo, amino, NHR, nitro, Me₃Sn, where R is as defined above or where the compound is a dimer with each monomer being the same or different and being of formula **Ia** or **Ib**, where the R₈ groups of the monomers form together a bridge having the formula -X-R¹-X- linking the monomers, where R¹ is an alkylene chain containing from 3 to 12 carbon atoms, which chain may be interrupted by one or more hetero-atoms and/or aromatic rings and may contain one or more carbon-carbon double or


Sub
01
triple bonds, and each X is independently selected from O, S, or N; or R₇ and R₈ together form a group -O-(CH₂)_p-O-, where p is 1 or 2; with the proviso that when A is a single bond, then R₂ is not CH=CR^AR^B, where R^A and R^B are independently selected from H, R^C, COR^C, CONH₂, CONHR^C, CONR^C, cyano or phosphonate, where R^C is an unsubstituted alkyl group having 1 to 4 carbon atoms.

Please amend claim 6 as follows:

6. (Twice amended.) A compound according to claim 1, wherein A is a single bond, and R₂ is an aryl group, or an alkyl or alkaryl group which contains at least one double bond which forms part of a conjugated system with a double bond of the pyrrolobenzodiazepine compound C-ring.

Please amend claim 13 as follows:

13. (Twice amended.) A compound of formula II:



wherein:

R₂' is O;


R₆, R₇ and R₉ are independently selected from H, R, OH, OR, halo, amino, NHR, nitro, Me₃Sn;

where R is a lower alkyl group having 1 to 10 carbon atoms, or an aralkyl group of up to 12 carbon atoms, whereof the alkyl group optionally contains one or more carbon-carbon double or triple bonds, which may form part of a conjugated system, or one or more carbonyl groups or one or more ether or thioether groups, or an aryl group of up to 12 carbon atoms; and is optionally substituted by one or more halo, hydroxy, amino, or nitro groups;

and where the compound is a dimer with each monomer being the same or different and being of formula II, where the R₈ groups of the monomers form together a bridge having the formula -X-R¹-X- linking the monomers, where R¹ is an alkylene chain containing from 3 to 12 carbon atoms, which chain may be interrupted by one or more hetero-atoms and/or aromatic rings and may contain one or more carbon-carbon double or triple bonds, and each X is independently selected from O, S, or N.

Please amend claim 20 as follows:

20. (Twice amended.) A compound of the formula **III**:



wherein:


CA
R₆, R₇ and R₉ are independently selected from H, R, OH, OR, halo, amino, NHR, nitro, Me₃Sn;

where R is a lower alkyl group having 1 to 10 carbon atoms, or an aralkyl group of up to 12 carbon atoms, whereof the alkyl group optionally contains one or more carbon-carbon double or triple bonds, which may form part of a conjugated system, or one or more carbonyl groups, or one or more ether or thioether groups, or an aryl group of up to 12 carbon atoms; and is optionally substituted by one or more halo, hydroxy, amino, or nitro groups;

and R₈ is amino.

Please amend claim 29 as follows:

29. (Once amended.) A compound of formula IV:



CS
wherein:

R₆, R₇ and R₉ are independently selected from H, R, OH, OR, halo, amino, NHR, nitro, Me₃Sn;

where R is a lower alkyl group having 1 to 10 carbon atoms, or an aralkyl group of up to 12 carbon atoms, whereof the alkyl group optionally contains one or more carbon-carbon double or triple bonds, which may form part of a conjugated system, or one or more carbonyl groups, or one or more ether or thioether groups, or an aryl group of up to 12 carbon atoms; and is optionally substituted by one or more halo, hydroxy, amino, or nitro groups;

R₈' and R₈'' are either independently selected from H, R or together form a cyclic amine; and

n is from 1 to 7.

Please amend claim 38 as follows:

38. (Twice amended.) A method of treating cancer comprising administering an effective amount of a compound according to claim 1, claim 13, claim 20 or claim 50 to a patient in need of such treatment wherein the cancer is selected from lung cancer, colon cancer, CNS cancer, melanoma, renal cancer, breast cancer and ovarian cancer.


CL
Sub
DI

Please amend claim 40 as follows:

Sub DI C7
40. (Twice amended.) A method of treating cancer comprising administering an effective amount of a compound according to claim 29 to a patient in need of such treatment wherein the cancer is selected from lung cancer, colon cancer, CNS cancer, melanoma, renal cancer and ovarian cancer.


Please amend claim 42 as follows:

42. (Once amended.) A process for preparing a compound according to claim 1 comprising cyclizing a compound of formula



wherein A, R₂, R₆, R₇, R₈ and R₉ are as defined in claim 1, R₁₀ is a nitrogen protecting group and CPQ is a masked aldehyde;


to a compound of formula



wherein A, R₂, R₆, R₇, R₈, R₉ and R₁₀ are as defined above and converting the above compound to a compound according to claim 1.


Please amend claim 43 as follows:

43. (Twice amended.) A method of treating a cisplatin-refractory disease comprising administering an effective amount to a patient in need of such treatment of a compound of formula



Please amend claim 44 as follows:

44. (Twice amended.) A method of inhibiting the growth of cisplatin-refractory cells which method comprises treating said cells with a compound of formula



Please amend claim 46 as follows:

- C11
Sub
D1
46. (Once amended.) A composition comprising a compound according to claim 1 and a pharmaceutically acceptable carrier or diluent.

Please amend claim 47 as follows:

- Sub
D1 C12
47. (Once amended.) A composition comprising a compound according to claim 13 and a pharmaceutically acceptable carrier or diluent.

Please amend claim 48 as follows:

- Sub
D1 C13
48. (Once amended.) A composition comprising a compound according to claim 20 and a pharmaceutically acceptable carrier or diluent.

Please amend claim 49 as follows:

- Sub
C14
49. (Once amended.) A composition comprising a compound according to claim 29 and a pharmaceutically acceptable carrier or diluent.

Please add the following claims:

56. (New) A compound according to claim 1 wherein


A is a single bond;

C/S
R₂ is 4-methoxyphenyl;

R₆ and R₉ are H, and


Sub
D1
R₇ and R₈ are methoxy.

57. (New.) A process for preparing a compound according to claim 13 comprising cyclizing a compound of formula




wherein R'₂, R₆, R₇, R₈ and R₉ are as defined in claim 13, R₁₀ is a nitrogen protecting group and CPQ is a masked aldehyde;

to a compound of formula




wherein R'₂, R₆, R₇, R₈, R₉ and R₁₀ are as defined above, and converting the above compound to a compound according to claim 13.

58. (New.) A process for preparing a compound according to claim 20 comprising cyclizing a compound of formula




wherein R₆, R₇, R₈, and R₉ are as defined in claim 20, R₁₀ is a nitrogen protecting group and CPQ is a masked aldehyde;
to a compound of formula




wherein R₆, R₇, R₈, R₉ and R₁₀ are as defined above, and converting the above compound to a compound according to claim 20.

59. (New.) A process for preparing a compound according to claim 29 comprising cyclizing a compound of formula



wherein R_6 , R_7 , R_8' , R_8'' , and R_9 are as defined in claim 29, R_{10} is a nitrogen protecting group and CPQ is a masked aldehyde;

to a compound of formula






wherein R_6 , R_7 , R_8' , R_8'' , R_9 and R_{10} are as defined above, and converting the above compound to a compound according to claim 29.

COC(=O)CC1=CNC(=O)c2cc(OC)c(BOC)c2N1CCOC(=O)CCc1c[nH]c2c1C(=O)c3cc(OC)c(OCC4=CC=CC=C4)cc3n2COc1cc(BOC)ccc1C(=O)N2C=CC(CO)C2COC(=O)CC1=CNC(=O)c2cc(OC)c(OC)cc2N1


C/S
Sub
D1

Sub
01


C/S



and




61. (New.) The method of claim 38 wherein the compound is



~~C15~~

62. (New.) The method of claim 38 wherein the compound is







Sub
01

CT5


63. (New.) The method of claim 40 wherein the compound is selected from the group consisting of

Sub
DI


C/S



and



64. (New.) The method of claim 38 wherein the compound is selected from the group consisting of



and

